

CERTAIN LOCALLY NILPOTENT VARIETIES OF GROUPS

ALIREZA ABDOLLAHI

ABSTRACT. Let $c \geq 0$, $d \geq 2$ be integers and $\mathcal{N}_c^{(d)}$ be the variety of groups in which every d -generator subgroup is nilpotent of class at most c . N.D. Gupta posed this question that for what values of c and d it is true that $\mathcal{N}_c^{(d)}$ is locally nilpotent? We prove that if $c \leq 2^d + 2^{d-1} - 3$ then the variety $\mathcal{N}_c^{(d)}$ is locally nilpotent and we reduce the question of Gupta about the periodic groups in $\mathcal{N}_c^{(d)}$ to the prime power finite exponent groups in this variety.

1. INTRODUCTION AND RESULTS

Let $c \geq 0$, $d \geq 2$ be integers and $\mathcal{N}_c$ be the variety of nilpotent groups of class at most c . We denote by $\mathcal{N}_c^{(d)}$ the variety of groups in which every d -generator subgroup is in $\mathcal{N}_c$. In [2], Gupta posed the following question:

For what values of c and d it is true that $\mathcal{N}_c^{(d)}$ is locally nilpotent?

Then he proved that for $c \leq (d^2 + 2d - 3)/4$, the variety $\mathcal{N}_c^{(d)}$ is locally nilpotent. In [1], Endimioni improved the latter result where he proved that for $c \leq 2^d - 2$, the variety $\mathcal{N}_c^{(d)}$ is locally nilpotent. Here we improve the number $2^d - 2$ to $2^d + 2^{d-1} - 3$. In fact we prove:

Theorem 1.1.

- (1) For $c \leq 2^d + 2^{d-1} - 3$, the variety $\mathcal{N}_c^{(d)}$ is locally nilpotent.
- (2) For $c \leq 2^d + 2^{d-1} + 2^{d-2} - 3$, every p -group in the variety $\mathcal{N}_c^{(d)}$ is locally nilpotent, where $p \in \{2, 3, 5\}$.

Note that the variety $\mathcal{N}_c^{(2)}$ is contained in the variety of c -Engel groups and it is yet unknown whether every c -Engel group is locally nilpotent, even, so far there is no published example of a non-locally nilpotent group in the variety $\mathcal{N}_c^{(2)}$. In the last section of this paper, by considering the problem of locally nilpotency of the variety $\mathcal{N}_c^{(2)}$, we study periodic groups in this variety. Note that since every two generator subgroup of a group in $\mathcal{N}_c^{(2)}$ is nilpotent, every periodic group in $\mathcal{N}_c^{(2)}$ is a direct product of p -groups (p prime). We reduce the question of Gupta for periodic groups in $\mathcal{N}_c^{(d)}$ to the locally nilpotency of p -groups of finite exponent in this variety where the exponent depends only on the numbers p and c . In fact we prove that

Theorem 1.2. Let p be a prime, $c > 1$ an integer and $r = r(c, p)$ be the integer such that $p^{r-1} < c - 1 \leq p^r$. Then the following are equivalent:

1991 *Mathematics Subject Classification.* 20F45.

Key words and phrases. Variety of groups, locally nilpotent groups .

This research was in part supported by a grant from IPM.

- (1) every p -group in $\mathcal{N}_c^{(2)}$ is locally nilpotent.
(2) if p is odd, every p -group of exponent dividing p^r in $\mathcal{N}_c^{(2)}$ is locally nilpotent, and if $p = 2$, every 2-group of exponent dividing 2^{r+1} in $\mathcal{N}_c^{(2)}$ is locally nilpotent.

2. GROUPS IN THE VARIETY $\mathcal{N}_c^{(d)}$

Let F_∞ be the free group of infinite countable rank on the set $\{x_1, x_2, \dots\}$, we define inductively the following words in F_∞ :

$$\begin{aligned} W_1 &= W_1(x_1, x_2) = [x_1, x_2, x_1, x_2], \\ W_n &= W_n(x_1, x_2, \dots, x_{n+1}) = [W_{n-1}, x_{n+1}, W_{n-1}, x_{n+1}] \quad n > 1; \\ V_1 &= V_1(x_1, x_2, x_3) = [[x_2, x_1, x_1, x_1, x_1], x_3, [x_2, x_1, x_1, x_1, x_1], x_3], \\ V_n &= V_n(x_1, x_2, x_3, \dots, x_{n+2}) = [V_{n-1}, x_{n+2}, V_{n-1}, x_{n+2}] \quad n > 1. \end{aligned}$$

For a group G and a subgroup H of G , we denote by $HP(G)$ the Hirsch-Plotkin radical of G and H^G the normal closure of H in G . We use the following result due to Heineken (see Lemma 8 of [3] and see Lemma 2 of [5] for the left-normed version).

Lemma 2.1. *Let G be a group and g an element in G such that $[g, x, g, x] = 1$ for all $x \in G$. Then the normal closure of $\langle g \rangle$ in $\langle g \rangle^G$ is abelian. In particular, $g \in HP(G)$.*

Lemma 2.2. *Let G be a group satisfying the law $W_n = 1$ for some integer $n \geq 1$. Then, G has a normal series $1 = G_n \triangleleft G_{n-1} \triangleleft \dots \triangleleft G_1 = G$ in which each factor G_i/G_{i+1} is locally nilpotent ($i = 1, 2, \dots, n-1$).*

Proof. We argue by induction on n . If $n = 1$, then Lemma 2.1 yields that $x_1 \in HP(G)$ for all $x_1 \in G$ and so G is locally nilpotent. Now suppose that the lemma is true for n and G satisfies the law $W_{n+1} = 1$. By Lemma 2.1, we have $W_n(x_1, \dots, x_{n+1}) \in HP(G)$ for all $x_1, \dots, x_{n+1} \in G$ and so $G/HP(G)$ satisfies the law $W_n = 1$. Thus by induction hypothesis $G/HP(G)$ has a normal series of length n with locally nilpotent factors, it completes the proof. $\square$

Lemma 2.3. *Let G be a p -group satisfying the law $V_n = 1$ for some integer $n \geq 1$ where $p \in \{2, 3, 5\}$. Thus G has a normal series $1 = G_{n+1} \triangleleft G_n \triangleleft \dots \triangleleft G_1 = G$ in which each factor G_i/G_{i+1} is locally nilpotent ($i = 1, 2, \dots, n$).*

Proof. We argue by induction on n . If $n = 1$, then by Lemma 2.1 $[x_2, x_1, x_1, x_1, x_1] \in HP(G)$ for all $x_1, x_2 \in G$. Thus $G/HP(G)$ is a 4-Engel group and since every 4-Engel p -group is locally nilpotent where $p \in \{2, 3, 5\}$ (see Traustason [9] and Vaughan-Lee [10]), so $G/HP(G)$ is locally nilpotent. Now suppose that G satisfies the law $V_{n+1} = 1$. By Lemma 2.1, $V_n(x_1, \dots, x_{n+2}) \in HP(G)$ for all $x_1, \dots, x_{n+2} \in G$, so $G/HP(G)$ satisfies the law $V_n = 1$. Thus by induction hypothesis, $G/HP(G)$ has a normal series of length $n+1$ with locally nilpotent factors, it completes the proof. $\square$

We use in the sequel the following special case of this well-known fact due to Plotkin [6] that every Engel radical group is locally nilpotent. (see also Lemma 2.2 of [1])

Lemma 2.4. *Let H be a normal subgroup of an Engel group G . If H and G/H are locally nilpotent, then G is locally nilpotent.*

Proof of Theorem 1.1. One can see that W_n and V_n are, respectively, in the $(2^{n+1} + 2^n - 2)$ th term and $(2^{n+2} + 2^{n+1} + 2^n - 2)$ th term of the lower central series of F_∞ , for all integers $n \geq 1$.

(1) every group G in the variety $\mathcal{N}_c^{(d)}$ satisfies the law $W_{d-1} = 1$ and so it follows from Lemmas 2.2 and 2.4, that G is locally nilpotent.

(2) if $d = 2$ then $c \leq 4$ and so every group in the variety $\mathcal{N}_c^{(d)}$ is 4-Engel. But as it is mentioned in the proof of Lemma 2.3, every 4-Engel p -group is locally nilpotent, where $p \in \{2, 3, 5\}$. Now assume that $d \geq 3$, then every group G in the variety $\mathcal{N}_c^{(d)}$ satisfies the law $V_{d-2} = 1$ and so it follows from Lemmas 2.3 and 2.4, that G is locally nilpotent. $\square$

3. p -GROUPS IN THE VARIETY $\mathcal{N}_c^{(2)}$

Lemma 3.1. *Let $c \geq 1$ be an integer, p a prime number and G a finite c -Engel p -group. Suppose that $x, y \in G$ such that $x^{p^n} = y^{p^n} = 1$ for some integer $n > 0$. Let r be the integer such that $p^{r-1} < c \leq p^r$. Then:*

- (a) *if p is odd and $n > r$ then $[x^{p^{n-1}}, y^{p^{n-1}}] = 1$.*
- (b) *if $p = 2$ and $n > r + 1$ then $[x^{2^{n-1}}, y^{2^{n-1}}] = 1$.*

Proof. Suppose that $K \leq H$ are two normal subgroups of G such that H/K is elementary abelian and a is an arbitrary element of G . Put $t = aK$ and $V = H/K$. Since $[V, {}_c t] = 1$, we have that $[V, {}_{p^r} t] = 1$ and $0 = (t - 1)^{p^r} = t^{p^r} - 1$ in $\text{End}(V)$. Thus $[H, a^{p^r}] \leq K$ for all $a \in G$. Now let $N = \langle x^{p^r}, y^{p^r} \rangle$. Then $[H, N] \leq K$ and since K, H are normal in G ; $[H, M] \leq K$ where $M = N^G$ the normal closure of N in G . Thus M is a normal subgroup of G centralizing every abelian normal section of G . By a result of Shalev [8]; if p is odd then M is powerful and if $p = 2$ then M^2 as well as all subgroups of M^2 which are normal in G , are powerful. Suppose that p is odd. Since M is generated by $\{(x^g)^{p^r}, (y^g)^{p^r} \mid g \in G\}$ and M is powerful, by Corollary 1.9 of [4], $M^{p^{n-r}}$ is generated by $\{(x^g)^{p^n}, (y^g)^{p^n} \mid g \in G\}$ and so $M^{p^{n-r}} = 1$. On the other hand $M^{p^{n-r-1}}$ is powerful by Corollary 1.2 of [4]. Thus $[M^{p^{n-r-1}}, M^{p^{n-r-1}}] \leq (M^{p^{n-r-1}})^p$. Now by Theorem 1.3 of [4], we have $(M^{p^{n-r-1}})^p = M^{p^{n-r}} = 1$. Thus $M^{p^{n-r-1}}$ is abelian and the part (a) has been proved. Now assume that $p = 2$. As it is mentioned in the above, since the subgroup $\langle x^{2^{r+1}}, y^{2^{r+1}} \rangle^G$ of M is normal in G , it is also powerful and the rest of the proof is similar to the latter case. $\square$

Lemma 3.2. *Let $c > 1$ be an integer, p a prime number and G a p -group in the variety $\mathcal{N}_c^{(2)}$ and let r be the integer satisfying $p^{r-1} < c - 1 \leq p^r$.*

- (a) *if p is odd then G^{p^r} is locally nilpotent.*
- (b) *if $p = 2$ then $G^{2^{r+1}}$ is locally nilpotent.*

Proof. Note that G is a c -Engel group. Suppose p is odd. First we prove that if x is an element of G such that $x^{p^n} = 1$ for some $n > r$, then $[a, x^{p^{n-1}}, x^{p^{n-1}}] = 1$ for all $a \in G$. Let $y = (x^{-1})^a$, then it is enough to show that $[y^{p^{n-1}}, x^{p^{n-1}}] = 1$. Since $\langle x, a \rangle \in \mathcal{N}_c$ then $\langle y, x \rangle \in \mathcal{N}_{c-1}$; therefore by Lemma 3.1, we have that $[y^{p^{n-1}}, x^{p^{n-1}}] = 1$. Now let z be an arbitrary element of G such that $z^{p^n} = 1$ for some positive integer n . We prove by induction on n that $z^{p^r} \in \text{HP}(G)$ and so it completes the proof for the case p odd. If $n \leq r$, then $z^{p^r} = 1 \in \text{HP}(G)$. Assume that $n > r$ then by induction hypothesis, $z^{p^{r+1}} \in \text{HP}(G)$. Now by the first part of the proof, $[a, z^{p^r}, z^{p^r}] = 1$

mod $HP(G)$, for all $a \in G$. So the normal closure of $z^{p^r}HP(G)$ in $G/HP(G)$ is abelian. Therefore $\langle z^{p^r} \rangle^G$ is (locally nilpotent)-by-abelian, hence Lemma 2.4 implies that $z^{p^r} \in HP(G)$.

The case $p = 2$ is similar. $\square$

Proof of Theorem 1.2. Suppose that p is odd and every p -group of exponent dividing p^r in $\mathcal{N}_c^{(2)}$ is locally nilpotent. Let G be a p -group in $\mathcal{N}_c^{(2)}$, then by Lemma 3.2(a), G^{p^r} is locally nilpotent. By assumption, G/G^{p^r} is locally nilpotent and since G is a c -Engel group, it follows from Lemma 2.4 that G is locally nilpotent. The case $p = 2$ is similar and the converse is obvious. $\square$

Now we use this result to some special cases. In fact we prove:

Proposition 3.3. *Every 2-group or 3-group in the variety $\mathcal{N}_5^{(2)}$ is locally nilpotent.*

Proof. By Theorem 1.2, we must prove that every 2-group of exponent dividing 8 and every 3-group of exponent dividing 9 in $\mathcal{N}_5^{(2)}$ is locally nilpotent. Suppose that G is a 2-group of exponent 8 in $\mathcal{N}_5^{(2)}$. Let $x, y \in G$, then $\langle x, y \rangle \in \mathcal{N}_5$ and of exponent 8. It is easy to see that $[x^4, y, x^4, y] = 1$. So by Lemma 2.1, $x^4 \in HP(G)$ for all $x \in G$. Therefore $G/HP(G)$ is of exponent dividing 4. By a famous result of Sanov (see 14.2.4 of [7]), $G/HP(G)$ is locally nilpotent and so by Lemma 2.4, G is locally nilpotent.

Now suppose that G is a 3-group of exponent 9 in the variety $\mathcal{N}_5^{(2)}$. Let $x, y \in G$, it is easy to see that $[x^3, y, x^3, y] = 1$. The rest of the proof is similar to latter case, but we may use this well-known result that every group of exponent 3 is nilpotent (see 12.3.5 and 12.3.6 of [7]) $\square$

REFERENCES

- [1] G. Endimioni, *Groups in which every d -generator subgroup is nilpotent of bounded class*, Quart. J. Math. Oxford (2), **46** (1995), 433-435.
- [2] N.D. Gupta, *Certain locally metanilpotent varieties of groups*, Arch. Math. **20** (1969), 481-484.
- [3] H. Heineken, *Engelsche Elemente der Länge drei*, Illinois J. Math. **5** (1961), 681-707.
- [4] A. Lubotzky and A. Mann, *Powerful p -groups. I. Finite groups*, J. Algebra **105** (1987), no. 2, 484-505.
- [5] M. L. Newell, *On right-Engel elements of length three*, Proc. Roy. Irish Acad. Sect. A **96** (1996), no. 1, 17-24.
- [6] B.I. Plotkin, *On the nil-radical of a group*, Dokl. Akad. Nauk SSSR (N.S.) **98** (1954), 341-343.
- [7] D.J.S. Robinson, "A first course in the group theory", Second edition, Graduate Texts in Mathematics, 80 Springer-Verlag, New York, 1996.
- [8] A. Shalev, *Characterization of p -adic analytic groups in terms of wreath products*, J. Algebra **145** (1992), no. 1, 204-208.
- [9] G. Traustason, *On 4-Engel groups*, J. Algebra **178** (1995), 414-429.
- [10] M. Vaughan-Lee, *Engel-4 groups of exponent 5*, Proc. London Math. Soc. (3) **74** (1997), no. 2, 306-334.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ISFAHAN; ISFAHAN 81746-73441; IRAN; AND INSTITUTE FOR STUDIES IN THEORETICAL PHYSICS AND MATHEMATICS, TEHRAN, IRAN.

E-mail address: a.abdollahi@sci.ui.ac.ir